

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 3/02, 25/00, 63/00, C05F 11/08	A1	(11) International Publication Number: WO 00/38513 (43) International Publication Date: 6 July 2000 (06.07.00)
(21) International Application Number: PCT/US99/29725 (22) International Filing Date: 14 December 1999 (14.12.99) (30) Priority Data: 09/222,459 29 December 1998 (29.12.98) US (71)(72) Applicant and Inventor: YAMASHITA, Thomas, T. [US/US]; 3631 Bogue Road, Denair, CA 95316-9619 (US). (74) Agent: FIELD, Bret, E.; Bozicevic, Field & Francis LLP, 200 Middlefield Road, Suite 200, Menlo Park, CA 94025 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: SOIL AMENDMENT COMPOSITIONS AND METHODS FOR USING THE SAME (57) Abstract Soil amendment compositions and methods for using the same are provided. The subject compositions are aqueous compositions consisting essentially of a carbon-skeleton energy component, a predisposing agent and a vitamin-cofactor component. The subject compositions find use in a variety of soil amendment applications, including: the control of soil born pests and pathogens; the improvement in soil fertility and/or characteristics, e.g. mineral release, water filtrations; the neutralization and/or degradation of toxins, etc.		

Document A
Cited in IDS for YAMA-008
Serial No. N/A,
filed Herewith

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SOIL AMENDMENT COMPOSITIONS AND METHODS FOR USING THE SAME

INTRODUCTION

5 Field of the Invention

The field of the invention is agriculture, particularly fertilizers.

Background

In agriculture, it is often desirable to use land that is not initially optimal in terms of soil fertility. Fertilizers have been developed for use on such land. Fertilizers are materials that
10 are used to supply elements needed for plant nutrition. Fertilizer materials may be in the form of solids, semi-solids, slurry suspensions, pure liquids, aqueous solutions and gases. Fertilizing materials may be introduced into a plant's environment in a number of different ways, including through addition to the soil, through application directly to a plant's foliage, and the like. The use of fertilizers is critical to commercial agriculture as fertilizers are
15 essential to correct natural deficiencies and/or replace components in soil. A number of different types of fertilizer compositions have been developed and employed in agriculture. However, there is continued interest in the development of new fertilizer compositions.

Another problem encountered by farmers and other agricultural workers is soil infestation with deleterious pests and/or pathogens. A variety of synthetic chemical pesticides
20 have been developed over the years to treat pest or pathogen infested soils. While such synthetic chemicals have been used with success, their use is not without controversy. Specifically, there is increasing public concern over the potential link between pesticide use and human disease conditions. As such, there is continued interest in the identification of new compositions which are capable of controlling soil borne pests or pathogens.

Relevant Literature

U.S. Patents of interest include: 5,549,729; 5,582,627; 5,696,094; 5,797,976; the disclosure of which are herein incorporated by reference.

5

SUMMARY OF THE INVENTION

Soil amendment compositions and methods for their use are provided. The subject compositions are aqueous compositions that include a predisposing agent, a carbon skeleton energy component and a vitamin-cofactor component. The subject compositions find use in a variety of different applications, including: the control of soil borne pests or pathogens; the neutralization and/or degradation of toxins; the improvement of soil characteristics, e.g. water permeability; the improvement of soil fertility; etc.

DESCRIPTION OF THE SPECIFIC EMBODIMENT

Aqueous compositions and methods for their use in soil amendment applications are provided. The subject compositions include a predisposing agent, a carbon-skeleton energy (CSE) source and a vitamin co-factor component. The subject compositions find use in a variety of different soil amendment applications, where such applications include: reducing the population of soil borne pests or pathogens; neutralizing or degrading soil toxins; improving soil characteristics; improving soil fertility; and the like. In further describing the invention, the compositions are described first in greater detail followed by a discussion of representative soil amendment methods in which the subject compositions find use.

Before the subject invention is described further, it is to be understood that the invention is not limited to the particular embodiments of the invention described below, as variations of the particular embodiments may be made and still fall within the scope of the appended claims. It is also to be understood that the terminology employed is for the purpose of describing particular embodiments, and is not intended to be limiting. Instead, the scope of the present invention will be established by the appended claims.

In this specification and the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs.

A critical component of the subject aqueous compositions is the predisposing agent. By predisposing agent is meant an agent which can traverse the membranes of a target pest or pathogen and thereby weaken the pest or pathogen physiology by virtue of denaturation of key enzymes (e.g. cytochrome oxidase) and/or proteins, and the like. A variety of different compounds are capable of fulfilling the above functional requirements and thereby serving as predisposing agents. As such, predisposing agents of interest include: aromatic amino acids, e.g. tyrosine, phenylalanine, tryptophan; phenols and derivatives and general products or reactants of the Shikimic Acid Pathway, e.g. cinnamic acid, chlorogenic acid, caffeic acid, coumaric acid, catechuic acid, ferulic acid, chorismic acid, quinic acid, gallic acid, gallotannins, scopoletin, dicoumarol, preocenes, phytoalexins such as orcinol, phaseolin, pisatin, isocoumarin, and the like; lignin alcohols e.g. coniferyl, sinapyl, p-coumaryl; flavonoids e.g. cyanidin, anthocyanidin, pelargonidin, delphinidin, malidin, peonidin, petunidin; flavonols and flavones; betalains e.g. betacyanin, betalain, betaxanthin; alkaloids e.g. caffeine, nicotine, theobromine; limonene -1-Methyl-4-(1-Methylethenyl) cyclohexene; p-mentha- 1, 8-diene; lignosulfonates, e.g. calcium lignosulfonate, potassium lignosulfonate, sodium lignosulfonate, ammonium lignosulfonate; and the like.

The subject compositions may include a single predisposing agent or a plurality of different predisposing agents. When a plurality of different predisposing agents are employed, the number of different predisposing agents may range from about 2 to 6, usually from about 2 to 5 and more usually from about 2 to 4.

The total amount of predisposing agent(s) that is present in the subject compositions is sufficient to achieve the desired weakening in the target pest or pathogen present in the soil. Generally, the amount present in the composition is sufficient to achieve a concentration in the soil following treatment ranging from about 5 to 650 ppm, usually from about 10 to 350 ppm and more usually from about 25 to 200 ppm. The amount of predisposing agent present in the composition may vary depending on whether the composition is to be applied "as is" to the soil or diluted prior to application, e.g. where the composition is a concentrate, as described infra. However, in many embodiments, the total amount of predisposing agent present in the composition ranges from about 5 to 75, usually from about 10 to 60 and more usually from about 15 to 60 % w/w of the composition.

A particularly preferred predisposing agent in many embodiments of the invention is a lignosulfonate. Lignosulfonates of interest include: calcium lignosulfonate, potassium

lignosulfonate, sodium lignosulfonate, ammonium lignosulfonate; and the like. The lignosulfonate is generally present in the composition in an amount that is sufficient to achieve a concentration in the soil following application that ranges from about 5 to 650 ppm, usually from about 10 to 350 ppm and more usually from about 25 to 200 ppm. In many
5 embodiments, the amount of lignosulfonate present in the composition ranges from about 5 to 75 % w/w, usually from about 10 to 50% w/w and more usually from about 15 to 60 % w/w of the composition. Lignosulfonates are generally available from Westway Terminal, Stockton CA; PM Ag, Stockton CA; Georgia Pacific, Bellingham WA; and the like.

Also of particular interest as a predisposing agent is gallic acid. When present, gallic
10 acid is present in an amount sufficient to achieve a soil concentration ranging from about .05 to 5.0 ppm, usually from about 0.1 to 5.0 ppm and more usually from about 0.2 to 1.0 ppm. In many embodiments, the amount of gallic acid present in the composition ranges from about 0.001 to 15 % w/w, usually from about 0.001 to 10% w/w and more usually from about .01 to 10% w/w.

15 In a particularly preferred embodiment, the subject soil amendment compositions include both a lignosulfonate and gallic acid. In such embodiments, the amounts of each of these agents in the composition are sufficient to achieve a lignosulfonate concentration in the treated soil that ranges from about 10 to 650 ppm and a gallic acid concentration in the soil that ranges from about 0.05 to 5.0 ppm. In many compositions falling within the scope of this
20 preferred embodiment, the amount of lignosulfonate ranges from about 10 to 50% w/w of the composition while the amount of gallic acid ranges from about 0.001 to 10% w/w of the composition.

A second critical component of the subject compositions is the carbon skeleton energy (CSE) component. CSE components that find use in the subject compositions are carbon
25 containing substances which provide a readily assimilable source of both carbon and energy for promoting microbial proliferation. Preferably, the CSE component provides a complex array of various carbon compounds such that varied enzymology is induced in microbes present in the target soil. As such, CSE sources that favor ancestral, beneficial species, which normally carry complex enzyme systems (as opposed to more simplified forms hosted by
30 facultative pathogens) are particularly preferred. Generally, the carbon skeleton energy component is a C₂ to C₁₀, usually C₄ to C₈ compound or polymer thereof, e.g. a polymer in which the monomeric units are C₂ to C₁₀ compounds, such as a polysaccharide. The CSE

component may be a single carbon containing compound or a composition of two or more different carbon containing or organic compounds. Compounds and compositions capable of serving as a CSE component include: complex organic compositions, such as molasses (e.g. cane, sugar beet, sorghum, etc.), whey, corn steep liquor, grape syrup, maple syrup, corn
5 syrup, etc; sugars, e.g. sucrose, fructose, glucose, lactose, galactose, dextrose, maltose, raffinose, ribose, ribulose, xylulose, xylose, amylose, arabinose, etc.; sugar phosphates, e.g. fucose-P, galactose-P, glucose-P, lactose-P, maltose-P, mannose-P, ribose-P, ribulose-P, xylose-P, xylulose-P, etc.; sugar alcohols, e.g. adonitol, sorbitol, mannitol, maltitol, ribitol, galactitol, glucitol, etc.; organic acids, e.g. glucuronic acid, alpha ketoglutaric acid,
10 galactonic acid, glucaric acid, gluconic acid, pyruvic acid, polygalacturonic acid, citric acid, succinic acid, malic acid, isocitric acid, folic acid, etc.; nucleotides and bases, e.g. adenosine, adenosine-P, uridine, uridine-P, thymine, thymine-P, cytosine, cytosine-P, guanine, guanine-P, etc.; and amino acids, e.g. glycine, alanine, leucine, isoleucine, asparagine, tyrosine, phenylalanine, serine, cysteine, valine, proline, methionine, glutamine, threonine, lysine,
15 aspartic acid, glutamic acid, arginine, and the like.

The CSE component is present in an amount sufficient to provide for a concentration in the target soil upon application that ranges from about 5 to 650 ppm, usually from about 10 to 350 ppm and more usually from about 25 to 200 ppm. In many embodiments, the amount of CSE component in the composition ranges from about 5 to 75 % w/w, usually from
20 about 10 to 50 % w/w and more usually from about 15 to 60 % w/w.

In a preferred embodiment, the CSE source is a molasses. Molasses may be obtained from a number of commercial sources, including cane molasses, etc., where commercial sources of molasses include: Westway Terminal, Stockton CA; PM Ag, Stockton, CA; and the like.

25 The final component is a vitamin-cofactor. A variety of agents are capable of serving as the vitamin-cofactor component of the subject aqueous compositions. Such agents include: yeast extract, yeast, vitamin Bs, e.g. thiamine pyrophosphate, riboflavin, biotin, pantothenic acid, phosphatidylcholine, inositol, PABA, nicotinic acid, folic acid and mixtures thereof, and the like. Of particular interest as a vitamin-cofactor is yeast extract, particularly yeast extract
30 obtained from spray dried extract, as available from Feedstuffs, Inc., Stockton CA; California Spray Dry, Stockton, CA; and the like. The amount of vitamin-cofactor component present in the composition is sufficient to provide a concentration in the treated soil that ranges from

about 0.01 to 10 ppm, usually from about 0.01 to 5.0 ppm and more usually from about 0.01 to 1 ppm. The amount of vitamin co-factor present in the composition generally ranges from about 0.001 to 15 % w/w, usually from about 0.001% to 10% w/w and more usually from about 0.01 to 5.0 % w/w.

5 Importantly, the composition includes substantially no macronutrients, e.g. nitrogen, potassium, phosphorous; or micronutrients, e.g. zinc, iron or manganese. By "substantially no" is meant that the amount of any one of these elements, if present in the composition, is insufficient for the element to influence the overall activity of the composition or the amendment of the target soil, i.e. the amount of the component is insufficient to make the
10 component an active ingredient of the composition. As such, for nitrogen, phosphorous and potassium, the amount present in the composition, if present at all, does not exceed about 5% and preferably does not exceed about 2.5%. For iron, zinc and manganese, the amount present in the composition, if present at all, does not exceed about 2% and preferably does not exceed about 1%. Despite the lack of these macro- and micronutrients, the composition
15 exhibits significant soil amendment activity.

 As the subject compositions are aqueous compositions, they further include a substantial amount of water. The amount of water present in the composition may vary depending on whether the composition is a concentrated or dilute composition. Generally, the compositions include at least about 5%, usually at least about 20% and more usually at least
20 about 30% water, where the amount of water present in the composition may be as high as 80% or higher, but generally does not exceed about 70% and usually does not exceed about 40%.

 The above soil amendment compositions are prepared by combining water with the various agents under conditions sufficient to produce an aqueous solution containing the
25 various agents. The water that is used to produce the subject compositions may be tap water obtained from any convenient water source, e.g. a municipal water district, where the water may be purified or otherwise treated, e.g. to remove certain undesirable agents that may be initially present therein. The various agents to be solubilized in the water to produce the soil amendment compositions may be obtained from any convenient source, e.g. commercial
30 vendor. For example, the carbohydrate component may be derived from a commercially available carbohydrate source, such as commercially available molasses, etc.

In preparing the subject soil fertilizer compositions, a concentrated or parent composition may first be produced, which parent composition or mix may or may not be diluted with water.

The subject aqueous compositions find use in a variety of soil amendment applications, i.e. methods of improving soil. In practicing the subject methods, the aqueous composition is contacted with the soil under conditions sufficient to achieve the desired concentrations of the agents of the composition in the soil. By contact is meant that the composition is introduced into the soil such that the desired concentration of the disparate components of the composition is obtained in the soil. As such, contact can include spraying so that the composition soaks into the soil, injecting the composition into the soil, flooding the soil with the composition, and the like. Contact is performed such that the concentration in the soil of the predisposing agent following treatment is at least about 5 ppm, usually at least about 20 ppm and more usually at least about 60 ppm, where the concentration of the predisposing agent following treatment may be as high as 650 ppm or higher, but generally does not exceed about 200 ppm and usually does not exceed about 60 ppm. Contacting also results in a concentration of the CSE component in the soil that is at least about 5 ppm, usually at least about 20 ppm and more usually at least about 60 ppm, where contact may result in a concentration of the CSE component that is 650 ppm or higher, but generally does not exceed about 200 ppm and usually does not exceed about 60 ppm. In addition, contact of the composition with the soil results in a vitamin-cofactor concentration in the soil that is at least about 0.01 ppm, usually at least about 0.05 ppm and more usually at least about 1.0 ppm, where the vitamin-cofactor concentration may be as high as 10 ppm or higher, but generally does not exceed about 5.0 ppm and usually does not exceed about 1.0 ppm.

The amount of aqueous composition that is used during any one application will vary greatly depending on the nature of the soil, the nature of the composition, the environmental conditions, etc. Where crops are treated with the subject compositions, the amount that is applied based on treated acreage is generally at least about 5 to 240 gal per acre, usually at least about 10 to 120 gal per acre, and more usually at least about 20 to 60 gal per acre, where the amount that is applied may be as high as 480 gal per acre or higher, but will usually not exceed about 240 gal per acre.

Depending on the nature of the soil, the nature of the composition, and the environmental conditions, as well as other factors, the composition may be applied more than

once over a given period of time. As such, the composition may be applied daily, weekly, every two weeks, monthly etc.

The aqueous compositions of the subject invention find use in a variety of different applications, where such applications include: the control of soil borne pests and pathogens; the improvement of water filtration; the improvement in mineral release; the enhancement in the water holding capacity of soil; the mellowing of soil textural qualities; the enhancement of the decomposition of plant tissues and accelerated degradation of potentially toxic chemicals and/or allelopathic chemicals; the improvement of root mass in plants grown in treated soil; and the like.

A variety of different soil borne pests may be controlled with the subject compositions. Such pests include: plant parasitic nematodes, phylloxera, grubs, and the like. By controlled is meant that the pest population in the soil is reduced, generally by at least about 5%, usually at least about 25% and more usually at least about 50%. As such, the invention provides methods and compositions for at least reducing, if not substantially eliminating, the population of soil borne pests in soil.

Similarly, the subject methods and compositions provide means for reducing the amount of pathogen present in soil. Pathogens that can be targeted with the subject methods include: pathogenic fungi, actinomycetes, bacteria, viruses, and the like. The subject methods result in a reduction of at least about 5 %, usually at least about 25%, and more usually at least about 50% of the amount of pathogen in the soil.

Also provided by the subject invention are methods and compositions for increasing indigenous soil microbe populations. Beneficial microbes whose population may be increased by the subject invention include: bacteria, fungi, actinomycetes, various free-living invertebrates, and the like. Applying the composition to the soil according to the subject methods results in at least a 2-fold increase, usually at least about a 20-fold increase and more usually at least about 40-fold increase in the microbe population in the treated soil.

The subject methods and compositions can also be used to improve water filtration through the soil. Water filtration may be improved by at least about 1.5×, usually at least about 2.5× and more usually at least about 4.5×.

Soil mineral release can also be enhanced using the subject methods and compositions. Mineral release, e.g. the release of minerals such as calcium, potassium and phosphorous, can

be improved by at least about 1.5×, usually at least about 3.0× and more usually at least about 5.0× as compared to that observed in control soil.

Finally, the subject methods and compositions can be used to increase the root mass of plants grown in the treated soil. Generally, the subject methods result in an increase in root mass of at least about 1.5×, usually at least about 2.0× and more usually at least about 4.0×, as compared to control plants, i.e. plants grown in untreated but otherwise substantially identical soil.

The following experiments are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

I. Tests To Indicate Efficacy of Various Predisposing Agents

15

The merits of alternative predisposing agents were examined by subjecting J2 larvae of the Root Knot Nematode (*Meloidogyne incognita*) to solutions of lignosulfonate and gallic acid. J2 larvae were exposed 24 hours to various concentrations of the materials in shallow 50 ml beakers. Following exposure, nematodes were inspected for activity level using touch response:

		Replications						Control %
Treatment		1	2	3	4	Total	Mean	
	1000 ppm Ligno	256/452	262/460	235/445	245/490	998/1847	250/462	45.9%
	5 ppm Gal Ac	305/462	316/440	326/447	302/450	1249/1799	312/450	30.7%
25	25 ppm Gal Ac	250/458	248/449	232/459	225/462	955/1828	239/457	47.7%
	100 ppm Gal Ac	193/450	185/448	176/436	150/464	704/1798	176/450	60.9%
	1000 ppm Gal Ac	25/449	39/444	40/454	27/460	131/1807	33/452	92.7%

Ligno =Ca Lignosulfonate

Gal Ac=Gallic Acid

Numbers listed are the active, responding nematodes per total examined (viable/total)

II. Compositions

Two amendment compositions were prepared: (a) an enhanced composition; and (b) a standard composition.

5

A. Enhanced (Enh) Composition Utilized

	<u>Component</u>	<u>Material Source</u>	<u>Amount % w/w</u>	<u>Final Concentration of a.i. in Total Mix</u>
10	Carbon Skeleton- Energy (CSE)	Hi- Brix Molasses	35%	17.5% CSE
	Predisposing Agent	Ca Lignosulfonate Gallic Acid	35% 0.10%	17.6% Predisp. Agent
15	Vitamin-Cofactor	Yeast Extract	2.5%	1.3 % Vit-Cofactor
	Water	Tap Water	27.5%	27.5 % Water
	Hi-Brix Molasses obtained from Westway Terminal, Stockton, CA			
	CaLignosulfonate obtained from Georgia-Pacific			
20	Gallic Acid obtained from Sigma Chemical Company			
	Yeast Extract obtained from Sigma Chemical Company			

B. Standard (Std) Composition Utilized

	<u>Component</u>	<u>Material Source</u>	<u>Amount % w/w</u>	<u>Final Concentration of a.i. in Total Mix</u>
25	CSE	Hi Brix Molasses	32%	16.0 % CSE
	Complexing Agent	Ca Lignosulfonate	32%	16.0 % Cplx
	Nitrogen	Urea (23%N)	5%	1.7 % Nitrogen
		KN ₃ (13.9%N)	3.8%	
30	Potassium	KN ₃ (38.7% K)	3.8%	1.5 % Potassium
	Phosphorus	H ₃ PO ₄ (23.7% P)	3.4%	0.8 % Phosphorus
	Zinc	ZnSO ₄ -7H ₂ O (36% Zn)	0.8%	0.3 % Zinc
	Iron	FeSO ₄ -7H ₂ O (31% Fe)	0.8%	0.3 % Iron
	Manganese	MnSO ₄ -H ₂ O (28% Mn)	0.8%	0.2 % Manganese
35	Vitamin B Cplx	Vitamin B Cplx	1.0%	0.04% B-Complex
	Water	Tap Water	20.4%	20.4 % Water

III. Comparative Examples

The above enhanced and standard compositions were compared in a variety of settings

5

A. *Enhancement of Indigenous Microbe Populations*

An inactive soil with low microbial activity was treated with 50 gpa Standard Mix and 50 gpa of the Enhanced Mixture and irrigated to incorporate the materials. Approximately 120 hours after treatment soil samples were secured and processed for examination of bacterial populations through dilution plating. Std and Enh are abbreviations for Standard and Enhanced Mix. The following results were observed:

Treatment	Replications					Total	Mean
	1	2	3	4	5		
Control	1.7	1.5	1.3	1.1	1.9	7.5	1.5 a
50 gpa Std	292.0	288.0	295.0	290.0	285.0	1,450.0	290.0 b
50 gpa Enh	605.0	595.0	598.0	610.0	592.0	3,000.0	600.0 c

Populations are expressed in millions of colony forming units per gram of soil

20 B. *Enhancement of Soil Mineral Release*

A relatively sterile soil was first examined for mineral release utilizing distilled water extraction. Portions of the field were then treated with 50 gpa or the standard mix (Std) and 50 gpa of the enhanced composition (Enh) and irrigated to maintain activation of the microbes. Six weeks following treatment, the soil were identically examined for mineral release. Model mineral elements examined were Ca, P and K. The following results were observed:

Treatment	<u>Calcium</u>			<u>Phosphorus</u>			<u>Potassium</u>		
	Before	After	Diff.	Before	After	Diff.	Before	After	Diff.
Control	987	1002	+15 ppm + 1.5%	8	8	0 0	45	47	+2 ppm +4.4%
5 50 gpa Std	992	1576	+584 ppm +58.9%	7	17	+10 ppm +143%	49	97	+48 ppm +98%
10 50 gpa Enh	972	2430	+1458 ppm +150%	8	32	+24 ppm +300%	43	150	+107ppm +249%

Except for %'s, all data are expressed as ppm mineral in the soil

C. Reductions In Disease Inoculum

- 15 Soil heavily infested with sclerotia of the pathogenic fungus, *Verticillium dahliae*, was examined for inoculum levels. Soils were then treated with 50 gpa Std and 50 gpa Enh material in which was mixed approximately 20 trillion colony forming units (cfu's) of beneficial bacteria and fungi, i.e. 1 gallon beneficial microbe suspension (IOTA, sold by FUSION 360, Stockton, CA). That is, 50 gpa Std material + 1 gallon of beneficial bacteria &
- 20 fungi were added per acre of soil. Likewise, 50 gpa Enh material + 1 gallon of beneficial bacteria & fungi were added per acre of soil. The soil was maintained near 80% field capacity to support microbial activity. Sixty days following the treatment the soils were again examined for inoculum levels of *V dahliae*. The following results were observed:

Treatment	Replications				T	M
	1	2	3	4		
25 Control Before	166	150	168	152	636	159
Control After	160	145	174	160	639	160
Difference	-6	-5	+6	+8	+3	+1 a
30 50 Std Bef	173	139	165	152	629	157
50 Std Aft	65	55	59	61	240	60
Difference	-108	-84	-106	-91	-389	-97 b
50 Enh Bef	180	172	166	170	688	172
35 50 Enh Aft	9	8	5	5	27	7
Difference	-171	-164	-161	-165	-651	-165 c

All cfu's of *Verticillium dahliae* are expressed per gram of soil

D. Reductions In Plant-Parasitic Nematode Populations

A naturally infested vineyard was initially evaluated for plant-parasitic nematode (PPN) populations. Vines were then treated with 50 gpa Std and 50 gpa Enh rates to which were each added 1 gallon of a bacterial-fungal suspension hosting more than 20 trillion cfu's per gallon. Nematode populations were again evaluated at 4 and 8 months. The following results were observed:

	Treatment	Replications				Total	Mean
		1	2	3	4		
10	Control Start	8,345	7,585	10,305	9,005	35,240	8,810
	Control 4 mos	11,025	7,225	8,455	12,985	39,690	9,923
	Control 8 mos	13,220	10,250	13,325	14,120	50,915	12,729
	Overall Change	+4,875	+2,665	+3,020	+5,115	+15,675	+3,919
15	50 Std Start	10,115	9,210	8,225	11,895	39,445	9,861
	50 Std 4 mos	6,125	5,350	4,035	3,980	19,490	4,873
	50 Std 8 mos	3,150	3,275	2,305	2,970	11,700	2,925 b
	Overall Change	-6,965	-5,935	-5,920	-8,925	-27,745	-6,936
20	50 Enh Start	12,845	9,205	7,995	12,430	42,475	10,619
	50 Enh 4 mos	345	415	200	190	1,150	288
	50 Enh 8 mos	75	20	15	nd	110	28 c
	Overall Change	-12,770	-9,185	-7,980	-12,430	-42,365	-10,591
25	All populations are the juvenile stages of Root-Knot Nematode, <i>Meloidogyne incognita</i> , per 250 cc of soil						

E. Increased Water Infiltration Rate

A soil with poor textural qualities was measured for water infiltration rate. The method of measurement involved use of a 4" diameter PVC pipe with one end shaped to facilitate each of placement into the soil. The PVC pipe was pushed into the soil to a depth of 4". A volume of water was then added to the cylinder and the rate of infiltration determined. Following initial measurements, the soil was treated with 50 gpa Std and 50 gpa of the Enh material + 1 gpa each of a bacterial-fungal suspension which contained more than 20 trillion cfu's per gallon. Material was sprayed evenly over the surface of the soil and gently incorporated with irrigation water. The soil moisture was maintained near 80% field capacity to encourage continuous microbial activity. All plots were again measured for rate of water infiltration 6 weeks following treatment. The following results were obtained:

Treatment	Replications			Total	Mean
	1	2	3		
Control Before	0.2	0.2	0.2	0.6	0.2
Control After	0.2	0.2	0.2	0.6	0.2
5 Overall Change	0	0	0	0	0 a
50 Std Before	0.2	0.2	0.2	0.6	0.2
50 Std After	0.6	0.6	0.6	1.8	0.6
10 Overall Change	+0.4	+0.4	+0.4	+1.2	+0.4 b
50 Enh Before	0.2	0.2	0.2	0.6	0.2
50 Enh After	0.8	0.8	0.8	2.4	0.8
15 Overall Change	+0.6	+0.6	+0.6	+1.8	+0.6 c

Values represent inches per hour water infiltration rate

F. *Increased Root Mass*

Bell pepper plants were treated with 50 gpa Std and 50 gpa Enh rates of the material. The treatment was supplemented with 1 gpa of a bacterial-fungal suspension hosting more than 20 trillion cfu's per gallon. Following 3 months of growth, the plants were gently uprooted and the root volume evaluated on a relative scale with the control assigned a value of 1.0. The following results were obtained:

Treatment	Replications					Total	Mean
	1	2	3	4	5		
25 Control	1.0	1.0	1.0	1.0	1.0	5.0	1.0 a
50 gpa Std	2.5	2.5	2.5	2.5	2.5	12.5	2.5 b
50 gpa Enh	3.4	3.2	3.0	3.2	3.4	16.2	3.2 c

Values represent relative root mass of bell peppers in relation to the control

It is evident from the above results and discussion that improved soil amendment compositions that are capable of improving soil in a number of different ways are provided. The compositions are relatively simple and easy to produce. Despite their simplicity, the compositions can provide for significant improvement in terms of soil characteristics, such as texture, water filtration, mineral release and the like. Furthermore, the compositions are useful in reducing the population of soil borne pests and pathogens in the soil. In addition, the compositions are made of natural products that do not pose a health risk to humans or livestock. As such, the subject compositions are a significant advance in the art.

All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

- 5 The invention now being fully described, it will be apparent to one of skill in the art that many changes and modifications can be made thereto without departing from the spirit and scope of the appended claims.

WHAT IS CLAIMED IS:

1. An aqueous composition consisting essentially of:
 - (a) a predisposing agent;
 - (b) a carbon-skeleton-energy component; and
 - 5 (c) a vitamin-cofactor.
2. The composition according to Claim 1, wherein said carbon-skeleton energy component comprises C₂ to C₁₀ molecules or polymers thereof.
- 10 3. The composition according to Claim 1, wherein said predisposing agent comprises a lignosulfonate.
4. The composition according to Claim 1, wherein said predisposing agent comprises gallic acid.
- 15 5. The composition according to Claim 1, wherein said predisposing agent comprises lignosulfonate and gallic acid.
6. An aqueous composition consisting essentially of:
 - 20 a carbon skeleton energy component;
 - a lignosulfonate;
 - gallic acid; and
 - a vitamin-cofactor.
- 25 7. The composition according to Claim 6, wherein said carbon-skeleton energy component is a molasses.
8. The composition according to Claim 6, wherein said vitamin-cofactor is yeast extract.
- 30 9. The composition according to Claim 6, wherein said carbon skeleton energy component is from about 10 to 50 % w/w of said composition.

10. The composition according to Claim 6, wherein said lignosulfonate is from about 10 to 50 % w/w of said composition.
11. The composition according to Claim 6, wherein said gallic acid is from about 0.001% to 10 % w/w of said composition.
12. The composition according to Claim 6, wherein said vitamin-cofactor is from about 0.001 to 10 % w/w of said composition.
13. An aqueous composition consisting of:
(a) from about 10 to 50 % w/w of a molasses;
(b) from about 10 to 50 % w/w of a lignosulfonate;
(c) from about 0.001 to 10 % w/w of gallic acid; and
(d) from about 0.001 to 10 % w/w of yeast extract.
14. The composition according to Claim 13, wherein said molasses makes up from about 30 to 40 % w/w of said composition.
15. The composition according to Claim 13, wherein said lignosulfonate makes up from about 30 to 40 % w/w of said composition.
16. The composition according to Claim 13, wherein said gallic acid makes up from about 0.01 to 1.0 % w/w of said composition.
17. The composition according to Claim 13, wherein said yeast extract makes up from about 2 to 3 % w/w of said composition.
18. The composition according to Claim 13, wherein water makes up from about 20 to 30 % w/w of said composition.

19. A method for amending soil, said method comprising:
applying to said soil an aqueous composition consisting essentially of:
- (a) a carbon-skeleton-energy component;
 - (b) a predisposing agent; and
 - 5 (c) a vitamin-cofactor.
20. The method according to Claim 19, wherein said method results in at least one of:
- (a) enhancement of the indigenous microbe population of said soil;
 - (b) enhancement of the mineral release ability of said soil;
 - 10 (c) reduction in disease inoculum present in said soil;
 - (d) reduction in parasitic nematode population in said soil;
 - (e) enhancement of water filtration through said soil; and
 - (f) enhancement in the fertility of said soil.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/29725

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 3/02, 25/00, 63/00; C05F 11/08

US CL : 71/1,11, 903; 424/405; 504/101, 113, 116, 118

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 71/1,11, 903; 424/405; 504/101, 113, 116, 118

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, Y	US 5,696,094 A (YAMASHITA) 09 December 1997 (09.12.97), see the abstract.	1-20
X, Y	US 5,549,729 A (YAMASHITA) 27 August 1996 (27.08.96), see the abstract.	1-20
X, Y	US 5,797,976 A (YAMASHITA) 25 August 1998 (25.08.98), see the abstract.	1-20
X, Y	US 5,582,627 A (YAMASHITA) 10 DECEMBER 1996 (10.12.96), see the claims and the abstract.	1-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

28 FEBRUARY 2000

Date of mailing of the international search report

19 APR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DEBORAH K. WARE

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/29725

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, USPATFULL, AGRICOLA, ANAABSTR, AQUASCI, BIOBUSINESS, BIOSIS, BIOCOMMERCE,
BIOTECHABS, BIOTECHDS, CABA, CANCERLIT, CAPLUS, CEABA, CEN, CIN, CONFSCI, CROPB, CROPU,
DDFB, DDFU, DGENE, DRUGB, DRUGLAUNCH, ETC.